

05 AGRICULTURE AND FORESTRY

Centre for Science and Environment
 41, Tughlakabad Institutional Area, New Delhi 110 062, INDIA
 Ph: +91-11-29956110 - 5124 - 6394- 6399 Fax: +91-11-29955879
 E-mail: cse@cseindia.org Website: www.cseindia.org

AGRICULTURE & FORESTRY




Emissions and mitigation potential

Agriculture, forestry and other land use (AFOLU) is a broad category of emissions that has been used by the IPCC since 2006. It widely used in national greenhouse gas inventories and combines two previously distinct sectors: Land use, land use change and forestry (LULUCF), and Agriculture

The AFOLU sector accounts for about a quarter (about 10-12 billion tons of CO₂ eq./year) of the net GHG emissions.¹ The main sources of emissions from the AFOLU sector comprise deforestation, agricultural emissions from soil and nutrient management and livestock. It is worth noting that uncertainties in calculation of net AFOLU emissions is larger than for other sectors; however some of the broad figures mentioned in the recent IPCC 5th assessment report are shown in the figure below:

Graph 1: Total anthropogenic GHG emissions (GtCO₂eq./yr) by economic sectors. Pull-out shows how indirect CO₂ emission shares (in % of total anthropogenic GHG emissions) from electricity and heat production are attributed to sectors of final energy use. Source: IPCC, 2014: Summary for Policymakers, In: Climate Change 2014, Mitigation of Climate Change.

Graph 1: Greenhouse gas emissions by economic sectors



Source: IPCC, 2014: Summary for Policymakers, In: Climate Change 2014, Mitigation of Climate Change.

Direct agricultural emissions


A wide range of agricultural activities emit GHGs and together they directly contributed about 11 per cent of total global anthropogenic emissions in 2010 (approximately 5 to 5.8 billion tons of CO₂ eq./year). Land-use and land-use change activities accounted for 9-11 per cent of total GHG emissions (4.3 to 5.5 billion tons of CO₂ eq./year).²

Emissions from energy use in agriculture added another 785 million tonnes of CO₂ eq. in 2010. The data includes emissions from fossil fuel energy needed to power machinery, irrigation pumps and fishing vessels.³

Most farm-related emissions come in the form of methane (CH₄) and nitrous oxide (N₂O). Enteric fermentation (belching and flatulence) from livestock and the addition of fertilisers and wastes to soils make up about 65 per cent of agriculture emissions globally.⁴ Other sources include rice cultivation, burning of crop residues and manure management.

Graph 2: Emissions from enteric fermentation were the greatest contributor to agricultural emissions (40 per cent), followed by manure left on pasture (16 per cent), synthetic fertilizers (13 per cent), rice cultivation

Graph 2: Emissions from enteric fermentation



Source: FN Tubiello et al., FAO working paper series, March 2014

Projected change in agricultural emissions in 2030 with respect to 2005

Type of emission	Total emissions 2005 (MtCO ₂ e) (million tonnes of CO ₂ equivalent)	Total emissions 2030 (projection) (MtCO ₂ e)	% change
Agricultural soils (N ₂ O)	1,840	2,483	35
Enteric fermentation (CH ₄)	1,894	2,320	22.5
Rice cultivation (CH ₄)	501	510	1.8
Manure management (CH ₄ , N ₂ O)	398	466	17
Other emissions (CH ₄ , N ₂ O)	1,165	1,165	0
Total non-CO ₂ emissions	5,798	6,944	20

(10 per cent), manure management (7 per cent) and burning of savanna (5 per cent)

Projections and trends for agriculture emissions

Total emissions from the agriculture sector (CH₄ and N₂O) are projected to increase by about 20 per cent by 2030. The largest increase will come from agricultural soils as can be seen in the table.⁵

Food supply chain emissions

Globally, roughly one-third of the food produced for human consumption, about 1.3 billion tonnes per year, is lost or wasted. Producing this much food accounts for 6-10 per cent of human-generated greenhouse gas emissions.⁶

- People in Europe and North America waste 95-115 kg of food each year, compared with only 6-11 kg a year for people in Sub-Saharan Africa and South Asia.⁷
- The average carbon footprint of food wastage is about 500 kg of carbon dioxide (CO₂) equivalents per person per year. Europe, North America, Oceania and industrial Asia have the highest per capita carbon footprint of food wastage

(approximately 700-900 kg of CO₂ equivalent per person per year), while sub-Saharan Africa has the smallest footprint per person (about 180 kg of CO₂ equivalents)⁸

- Developing countries have high losses at the post-harvest and processing stages because of spoilage. Spoilage is high because of lack of modern transport and storage infrastructure, and financial, managerial and technical limitations in difficult climatic conditions.⁹
- The total amount of food wasted by consumers in industrialised countries is nearly as high as the total net food production in sub-Saharan Africa.¹⁰
- A total of 1.4 billion ha of land – 28 per cent of the world's agricultural area – is used annually to produce food that is lost or wasted. This is equivalent to the area of Canada and India put together, and dwarfed globally only by the size of Russia (1.7 billion ha).¹¹
- Food wastage costs food producers some USD 750 billion annually.¹²
- High-income regions and Latin America are responsible for about 80 per cent of all meat waste.¹³

Graph 3: About a third of all food produced is lost in the food supply chain



Source: CGIAR Big Facts: <http://ccafs.cgiar.org/bigfacts2014>

Emissions from livestock

- The livestock sector contributes an estimated 7,100 million tonnes of CO₂ equivalent per year, representing 14.5 per cent of human-induced greenhouse gas emissions.¹⁴
- Production of beef and cattle milk accounts for the majority of emissions, contributing 41 per cent and 20 per cent respectively of the sector's emissions. The production of pig meat and poultry meat and eggs contribute 9 per cent and 8 per cent respectively of the sector's emissions.¹⁴
- Feed production and processing, and enteric fermentation from ruminants are the two main sources of emissions, accounting for 45 per cent and 39 per cent respectively of sector emissions. Manure storage and processing account for 10 per cent of emissions.¹⁴
- The consumption of fossil fuel along livestock-sector supply chains accounts for about 20 per cent of sector emissions.¹⁴
- About 44 per cent of the sector's emissions are in the form of methane (CH₄). The rest is almost equally shared between N₂O (29 per cent) and CO₂ (27 per cent).¹⁴
- Global CH₄ emissions from enteric fermentation increased by 7 per cent from 1990 to 2005, from 1,763 million tonnes of CO₂ equivalent to 1,894 million tonnes, corresponding to 32.7 per cent of total agricultural emissions.¹⁵
- From 2005 to 2030, CH₄ emissions from enteric fermentation are projected to increase by 22 per cent, from 1,894 million tonnes of CO₂ equivalent to 2,320 million tonnes.¹⁵
- From 2005 to 2030, global CH₄ and N₂O emissions from manure management are projected to increase by 17 per cent, from 398 million tonnes of CO₂ equivalent to 466 million tonnes (US-EPA 2012 pp. 72-73).¹⁵

Equity in agriculture and livestock emissions

Agriculture has to be put at the heart of any poverty-reduction strategy. It is a multi-dimensional sector directly linked to the fight against hunger and malnutrition and to food security. At the same time, it is strongly influenced by international trade, finance, development cooperation and, increasingly, it is affected by climate change and environmental degradation. This diversity of functions and activities requires that issues relating to agriculture be treated in a holistic manner, and that the challenge of policy coherence be tackled both at national and international levels.¹⁶

India, China and a large number of African countries have countered by pointing out that emission-reduction efforts in the agricultural sector would affect farmers, who constitute a large percentage of the population and are often the poorest, in the developing world.¹⁷ They also argue that the effort to reduce emissions should focus on fossil-fuel-based activities that spew out carbon dioxide – the greatest contributor to global warming by far. As paddy fields and livestock are some of the biggest causes of emissions, emission reduction in the sector has major implications for India and China.

Land-use change emissions and mitigation potential of forests

- Land-use change is a highly complex process. It results from the interaction of diverse drivers that may be direct or indirect and can involve numerous transitions, such as clearing, grazing, cultivation, abandonment and secondary regrowth.²⁰
- Deforestation and land-use change accounts for 2,200-6,600 million tonnes of carbon dioxide (CO₂) equivalent per year, or 30-50 per cent of agricultural emissions and about 4-14 per cent of global emissions.²¹
- Land-use change to produce feed for livestock

Agriculture emissions – developed versus developing countries¹⁸

Developed countries, with 17 per cent of the world population, are responsible for 26 per cent of global N₂O emissions from soils, 30 per cent of methane CH₄ emissions from enteric fermentation, and 52 per cent of CH₄ and N₂O emissions from manure management. Agriculture emissions from developed countries increased by nearly 17 per cent between 1990 and 2005 due in good part to a massive increase in nitrogen fertiliser use in New Zealand and Australia and an increase of manure effluent of cattle, poultry and swine farms and manure application to soils in North America.

Despite roughly similar numbers of cattle per capita and many more small ruminants per capita in Africa than in developed countries, the latter's methane (CH₄) and nitrous oxide (N₂O) emissions from livestock manure are over 10 times greater than from African countries. Why? Because confined industrial animal agriculture stores manure in immense lagoons that produce significant quantities of these greenhouse gases.


Illustrative comparison of developed and developing country agriculture emissions

Country	Non-CO ₂ GHG emissions from agriculture in 2010 (Mt CO ₂ eq)	Per capita non-CO ₂ emissions from agriculture (tonnes CO ₂ eq)	Per capita N ₂ O emissions from agricultural soils (tonnes CO ₂ eq)	Per capita emissions from livestock (from livestock) (tonnes CO ₂ eq)	Per capita meat consumption (kg)
France	102.7	1.63	0.86	0.77	88.6
Ghana	8.7	0.36	0.07	0.08	9.9
India	732.3	0.60	0.18	0.18	5.2
Malawi	12.8	0.86	0.66	0.09	5.1
New Zealand	37.1	8.49	2.91	5.59	142.1
United States	454.9	1.46	0.78	0.64	124.8


Source: Climate-Justice 2012¹⁹

Agriculture is responsible for 75 per cent of global deforestation

If trends continue, approx **10 million km²** of land will likely be cleared by 2050 to meet food demand



Alternative pathways would only require approx **2 million km²** of land be cleared



Source: CGIAR Big Facts: <http://ccafs.cgiar.org/bigfacts2014/#>

accounts for about 9 per cent of livestock-sector emissions.²²

- Since 1850, land-use change has directly contributed some 35 per cent of human-generated CO₂ emissions.²³
- Past trends imply that about 10 million square km of land will be cleared by 2050 to meet food demand, leading to annual emissions of 3,000 million tonnes of CO₂ equivalent. Reducing land clearing to about 2 million square km would reduce greenhouse gas emissions to 1,000 million tonnes of CO₂ equivalent per year.²⁴

Co-benefits of mitigation in agriculture

Temperature rise above 2°C will produce increasingly unpredictable and dangerous impacts for people and ecosystems, but particularly for agricultural systems. Impacts on the agricultural sector that are already

occurring but expected to intensify include increased irrigation water needs, increased spread of animal and crop diseases and pests, reduced forage quality, and reduced crop and pasture yields. These impacts stem from changes in surface temperatures, timing of seasons and frequency and severity of severe weather events, such as droughts, floods, and heatwaves.²⁵

AFOLU plays a central role for food security and sustainable development. The most cost-effective mitigation options in forestry are afforestation, sustainable forest management and reducing deforestation, with large differences in their relative importance across regions. In agriculture, the most cost-effective mitigation options are cropland management, grazing land management, and restoration of organic soils.

The main mitigation options within AFOLU, as listed in the IPCC 5th Assessment Report chapter 11 on Agriculture, involve one or more of these strategies:

- reduction/prevention of emissions to the atmosphere by conserving existing carbon pools in soils or vegetation that would otherwise be lost or by reducing emissions of CH₄ and N₂O
- sequestration – enhancing the uptake of carbon in terrestrial reservoirs, and thereby removing CO₂ from the atmosphere
- reducing CO₂ emissions by substitution of biological products for fossil fuels or energy intensive products
- Demand side options (e.g. by lifestyle changes, reducing losses and wastes of food, changes in human diet, changes in wood consumption), though known to be difficult to implement, may also play a role.

The IPCC 5th Assessment Report also estimates the potentials for reducing greenhouse gases in agriculture as follows:

- Global economic mitigation potentials in agriculture in 2030 are estimated to be 0.5-10.6 GtCO₂eq/yr
- Reducing food losses and waste can reduce GHG emissions by 0.6-6.0 GtCO₂eq/yr
- Changes in diet could result in GHG emission savings of 0.7-7.3 GtCO₂eq/yr
- Forestry mitigation options are estimated to contribute 0.2-13.8 GtCO₂/yr

Note: The points under Food Supply, Livestock and Forestry emissions were adapted from the CGIAR Big facts an open-access resource. <http://ccafs.cgiar.org/bigfacts2014/#about=true>

References

1. IPCC, 2014: Climate Change 2014: Mitigation of Climate Change. Chapter 11 Agriculture, Forestry and Other Land Use (AFOLU).
2. Ibid.
3. FAO, 2014. <http://www.fao.org/resources/infographics/infographics-details/en/c/218650/>
4. <http://www.ipcc-nggip.iges.or.jp/public/2006gl/>
5. CGIAR Big Facts: <http://ccafs.cgiar.org/bigfacts2014/#>
6. Vermeulen et al. 2012. Climate change and food systems. *Annual Review of Environmental Resources* 37:195–222
7. Gustavsson et al. 2011. Global food losses and food waste. <http://www.fao.org/docrep/014/mb060e/mb060e00.pdf>
8. Food and Agriculture Organization of the United Nations. 2013a. Food wastage footprint. Impacts on natural resources. Summary report.
9. Gustavsson et al. 2011. Global food losses and food waste. <http://www.fao.org/docrep/014/mb060e/mb060e00.pdf>
10. Food and Agriculture Organization of the United Nations. 2013a. Food wastage footprint. Impacts on natural resources. Summary report
11. Ibid.
12. Ibid.
13. Ibid.
14. Gerber et al. 2013. Tackling climate change through livestock – A global assessment of emissions and mitigation opportunities. FAO.
15. United States Environmental Protection Agency. 2012. Global anthropogenic non-CO₂ greenhouse gas emissions: 1990 – 2030.
16. UNCTAD 2013. Trade and Environment Review 2013. Wake up before it is too late: Make agriculture truly sustainable now for food security in a changing climate
17. <http://www.thehindu.com/sci-tech/energy-and-environment/india-scores-a-win-in-warsaw-on-emission-cuts-affecting-farmers/article5347727.ece>
18. UNECA 2011. Climate Change Impacts on Agriculture in Africa and the UNFCCC negotiations: Policy implication of recent scientific findings. <http://climate-justice.info/wp-content/uploads/2011/12/Climate-and-agriculture-AGN-paper.pdf>

19. <http://climate-justice.info/wp-content/uploads/2012/11/Equity-and-Ambition-in-Agriculture.pdf>
20. Gerber et al. 2013. Tackling climate change through livestock – A global assessment of emissions and mitigation opportunities. FAO
21. Vermeulen et al. 2012. Climate change and food systems. *Annual Review of Environmental Resources* 37:195–222
22. Gerber et al. 2013. Tackling climate change through livestock – A global assessment of emissions and mitigation opportunities. FAO
23. Foley et al. 2005. Global consequences of land use. *Science* 309:570–574
24. Tilman et al. 2011. Global food demand and the sustainable intensification of agriculture. *PNAS* 108(50):20260–20264. (Available from www.pnas.org/cgi/doi/10.1073/pnas.1116437108)
25. GHG protocol Agriculture guidance: <http://www.ghgprotocol.org/standards/agriculture-guidance>